

SUNOSI improves wakefulness in adults with excessive daytime sleepiness due to obstructive sleep apnea (OSA) or narcolepsy. SUNOSI does not treat the cause of OSA or take the place of your CPAP. Continue to use any treatments or devices as prescribed by your doctor.



UNABLE TO STAY AWAKE DURING THE DAY? TALK TO YOUR DOCTOR

Excessive daytime sleepiness (EDS) is a medical condition that can be caused by obstructive sleep apnea (OSA) or narcolepsy. With EDS, you may have an irresistible need to sleep during the day and feel tired all the time.

The first step in finding a solution is talking with your doctor. The following topics can help get the conversation started.

I have

- Not been diagnosed with EDS, but extreme daytime tiredness is a consistent part of my life
- Been diagnosed with EDS due to
 - OSA
 - Narcolepsy

Daytime sleepiness has been affecting me for

- Less than 1 month
- 1 to 6 months
- 6 months to 1 year
- 1 to 2 years
- More than 2 years

I currently use CPAP to treat my OSA

- Yes
- No
- I used CPAP, but don't any longer
- Not applicable

I am currently treating my EDS with medication

- Yes
- No
- I was prescribed medicine, but am no longer taking it
- Not applicable

The medicine I am taking for my EDS is

- Not working well for me
- Working well for me
- I am not on treatment for EDS

My excessive daytime sleepiness impacts my life in the following ways

- It negatively impacts my relationships with family, friends, and coworkers
- I regularly avoid activities with friends because I'm so tired
- I fall asleep or doze off while stopped in traffic
- I fall asleep at work when I should not
- I fall asleep regularly while relaxing (e.g., watching TV)
- Other _____

If there are other questions you want to ask, or details you need to share, note them here

During your appointment, ask your doctor

- Are there tests, like the Epworth Sleepiness Scale, that can help assess my level of sleepiness?

Talk to your doctor about your answers and how extreme daytime tiredness is affecting you.

INDICATION AND IMPORTANT SAFETY INFORMATION

What is SUNOSI? SUNOSI (solriamfetol) is a prescription medicine used to improve wakefulness in adults with excessive daytime sleepiness due to narcolepsy or obstructive sleep apnea (OSA).

- SUNOSI does not treat the underlying cause of OSA and SUNOSI does not take the place of any device prescribed for OSA, such as a continuous positive airway pressure (CPAP) machine. It is important that you continue to use these treatments as prescribed by your healthcare provider.

IMPORTANT SAFETY INFORMATION

Do not take SUNOSI if you are taking, or have stopped taking within the past 14 days, a medicine used to treat depression called a monoamine oxidase inhibitor (MAOI).

Please see additional Important Safety Information on the next page and Medication Guide.

IMPORTANT SAFETY INFORMATION (CONT'D)

Before taking SUNOSI, tell your doctor about all of your medical conditions, including if you:

- have heart problems, high blood pressure, kidney problems, diabetes, or high cholesterol.
- have had a heart attack or a stroke.
- have a history of mental health problems (including psychosis and bipolar disorders), or of drug or alcohol abuse or addiction.
- are pregnant or planning to become pregnant. It is not known if SUNOSI will harm your unborn baby.
- are breastfeeding or plan to breastfeed. SUNOSI passes into your breast milk. Talk to your doctor about the best way to feed your baby if you take SUNOSI.

What are the possible side effects of SUNOSI?

SUNOSI may cause serious side effects, including:

Increased blood pressure and heart rate. SUNOSI can cause blood pressure and heart rate increases that can increase the risk of heart attack, stroke, heart failure, and death. Your doctor should check your blood pressure before, and during, treatment with SUNOSI. Your doctor may decrease your dose or tell you to stop taking SUNOSI if you develop high blood pressure that does not go away during treatment with SUNOSI.

Mental (psychiatric) symptoms including anxiety, problems sleeping (insomnia), irritability, and agitation. Tell your doctor if you develop any of these symptoms. Your doctor may change your dose or tell you to stop taking SUNOSI if you develop side effects during treatment with SUNOSI.

The most common side effects of SUNOSI include:

- headache
- decreased appetite
- problems sleeping
- nausea
- anxiety

These are not all the possible side effects of SUNOSI. Call your doctor for advice about side effects.

SUNOSI (solriamfetol) is available in 75 mg and 150 mg tablets and is a federally controlled substance (CIV) because it contains solriamfetol that can be a target for people who abuse prescription medicines or street drugs. Keep SUNOSI in a safe place to protect it from theft. Never give or sell your SUNOSI to anyone else because it may cause death or harm them and it is against the law. Tell your doctor if you have ever abused or been dependent on alcohol, prescription medicines, or street drugs.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call **1-800-FDA-1088**.

Please see [Medication Guide](#).

SUN CON ISI 06/2023

For more information, visit SUNOSI.com.

axsome

SUNOSI, AXSOME, and its logos are trademarks or registered trademarks of Axsome Therapeutics, Inc. or its affiliates. For U.S. residents ages 18 and older.
© 2023 Axsome Therapeutics, Inc. All rights reserved.

One World Trade Center, 22nd Floor, New York, NY 10007. PP-SUN-US-2200263 08/2023

sunosI
(solriamfetol) 
75, 150 mg tablets